

A MATEMATIKA ALAPJAI, 8. ELŐADÁS

András Kornai

BMETE91AM35 2021-22 Őszi félév

- Kétféle igazságfogalom van: szintaktikai és szemantikai
- Jelben \vdash ‘-ből következik’ or -ből levezethető’ ahol $A \vdash B$ azt jelenti B formálisan levezethető (bizonyítható) A -ból. Pl. a legtöbb rendszerben $x = 3 \wedge y = x \vdash y = 3$, és van egyfajta *bizonyításelmélet* (proof theory) ami képes erre. Ebben csak képleteket manipulálunk: meglévő képletekből újakat vezetünk le mechanikusan
- Van továbbá a \models ‘modellálja’ reláció ahol $A \models B$ azt jelenti, hogy minden modellben ahol A teljesül ott B is igaz. Ehhez *modellelemélet* kell ami megmondja, hogy mi a viszony egy elmélet (formulahalmaz) és struktúrált halmazok (modellek) közt
- Jól megcsinált rendszerekben $A \vdash B$ -ből következik $A \models B$

A FORDÍTOTTJA NEM IGAZ!

- In many well-crafted systems (e.g. the first order formulation of Peano Arithmetic) there are statements which are semantically true e.g. $PA \models \text{Goodstein's Theorem}$, but *has no proof there*
- If it has no proof, how do we know it's true? Because in a stronger system (in this case, 2nd order arithmetic) we can prove it
- That the converse is not true for systems endowed with a bit of arithmetic is the celebrated Gödel Incompleteness Theorem
- Our interest here is with the less celebrated, but just as important, Gödel Completeness Theorem
- This says that every formula that is true in all structures is provable
- Wait, how can these both be true? The answer is that PA has more models in first-order axiomatization than in second-order

GOODSTEIN'S THEOREM

- When we write a natural number n in base b this means we express it as $\sum_{i=0}^k c_{j_i} b^{e_{j_i}}$ where the digits c_{j_i} are between 1 and $b - 1$ and the exponents e_{j_i} are arbitrary nonnegative integers.
- In *hereditary* base b notation we force the exponents to be also written in base b and the exponents therein, and so on, until all digits are $< b$ everywhere.
- The *Goodstein sequence* $G(n)(k)$ for a number n begins with $G(n)(2) = n$, written in hereditary base 2. Next replace each occurrence of 2 by 3 (in general, each occurrence of b by $b + 1$) and subtract 1. Since you increase the base and subtract only one the sequence increases very very fast.
- **Goodstein's Theorem** Every Goodstein sequence terminates in 0 in finitely many steps
- Once you have the theory of *ordinals* at your disposal, the theorem is easy to prove. But no proof without such powerful theory exists (Kirby and Paris 1982).

PROPOSITIONAL LOGIC

- Any statement that can be true or false (in either of the senses discussed above) is called a **proposition**. These come in two basic varieties: *a has property P* and *the relation R holds between some elements*. Examples of the first: *57 is prime*, of the second: $2 + 3 = 4$
- Things that are *not* propositions include imperatives *Go home!* and questions *Where is Johnny?*
- Declarative statements using variables are called **open propositions** 'x is prime'. These can get a truth value either by substitution '17 is prime' and '18 is prime' both have a truth value or by quantification (part of FOL, but not PL)
- ZFC Axiom 3 (comprehension) creates the connection between PL and set theory: for any open sentence $\phi(x, w_1, \dots, w_n)$ and any set A there exists a set B containing all and only those elements x of A for which $\phi(x, w_1, \dots, w_n)$ holds. 'elements of a set satisfying some proposition can be collected in a set'

BOOLEAN OPERATIONS

- Well, what are operations? Operations are like addition, multiplication, negation... How can we define operations?
- We don't need new machinery! *Binary* operations are **functions** with two variables. *Unary* operations are functions with one variable (minus, reciprocal, ...) *Nullary operations* are functions that don't depend on any variable, **constants**.
- A **structure** is a set S and some operations. For example groups have a nullary operation (the unit e), a unary operation ($^{-1}$), and a binary operation (multiplication) which satisfy some identities (group axioms). On occasion, we don't insist that an operation be everywhere defined.
- One set of operations that matters in PL are the Boolean \neg, \wedge, \vee
- These are 'truth functional' – only the truth of the operands matters for establishing the truth of the result

IMPLICATION IN PL

- 1 We define $P \rightarrow Q$ by $\neg P \vee Q$
- 2 This has interesting consequences, the most important being *ex falso quidlibet* 'everything follows from a false statement' or *if you start with a false premiss, you can derive any conclusion from it*
- 3 We also define truth-functional equivalence $P \equiv Q$ by $P \rightarrow Q \wedge Q \rightarrow P$
- 4 Remember sets satisfied the de Morgan identities $\overline{A \cup B} = \overline{A} \cap \overline{B}$ and $\overline{A \cap B} = \overline{A} \cup \overline{B}$? In PL they are satisfied by $\neg(A \vee B) = \neg A \wedge \neg B$ and $\neg(A \wedge B) = \neg A \vee \neg B$

HÁZI FELADATOK

- Jövő hétre: CPZ 2.8; 2.26; 2.32; 2.44; 2.59 és a következő HW8.6: Ismerd meg a latex tabular környezetet és készíts igazságtáblát a következő oszlopokkal:
 $P, Q, \neg P, \neg Q, P \rightarrow Q, \neg Q \rightarrow \neg P$
- Csak a pdf kell (a latex forráskód nem) és az email tárgya
Subject: MATALAP NEPTUN HW8